



RESEARCH ARTICLE

Studies on Compatibility of *Metarhizium anisopliae* with Insecticides

Kiruthiga G^{1*}, Jeyarani S¹, Sathiah N¹, Murugan M¹, Sivakumar U² and Uma D³

^{1*}Department of Agricultural Entomology, Tamil Nadu Agricultural University, Coimbatore-641 003, India

²Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore-641 003, India

³Department of Agricultural Biochemistry, Tamil Nadu Agricultural University, Coimbatore-641 003, India

ABSTRACT

The results on the compatibility of *M. anisopliae* (isolate TNAU-MA-GDU) with various insecticides at five different doses (0.1x, x, 2x, 5x, and 10x) revealed that the highest mycelial growth was observed in flubendamide 480 SC (89.00 to 78.8 mm) and spinetoram 11.70 SC (89.00 to 76.60 mm) and was significantly superior over rest of insecticides tested. The minimum growth of *M. anisopliae* was observed in insecticidal treatment with azadirachtin (Neem 1500) (65.2 to 18.2 mm) followed by novaluron 10 EC (86.6 to 42.1 mm). All the tested insecticides except azadirachtin (Neem 1500) were compatible with *M. anisopliae* (TNAU-MA-GDU) at lower and field doses. The results on the compatibility of insecticides with *M. anisopliae* clearly indicate that the pesticides having better compatibility exhibited the maximum mycelial growth, sporulation, and biomass production in order of flubendamide 480 SC > spinetoram 11.70 SC > emamectin benzoate 5 SG > chlorantraniliprole 18.5 SC > novaluron 10 EC > azadirachtin (Neem 1500).

Received: 12 July 2022

Revised: 12 August 2022

Accepted: 11 September 2022

Keywords: Compatibility; Vegetative growth; T value; *Metarhizium anisopliae*; and Insecticides

INTRODUCTION

Insecticides have different effects on the fungal growth, sporulation, and germination of entomopathogenic fungi depending on the fungal species (Schumacher and Poehling, 2012). The combination of low doses of insecticides and entomopathogenic fungi is an added advantage in IPM practices as it minimises the insecticides use, reduces environmental pollution, and weakens resistance to insect pests (Sain *et al.*, 2019). Effective control of insect pests can be achieved by using a combination of entomopathogenic fungi and insecticides (Oliveira *et al.*, 2003; Purwar and Sachan, 2006; Asi *et al.*, 2010). Conversely, the usage of non-selective or incompatible insecticides with entomopathogenic fungus may potentially prevent fungal growth and development, resulting in the failure of IPM. The knowledge of the compatibility between entomopathogenic fungus and pesticides is essential to choose appropriate chemicals and to decide on the timing of treatments to prevent the harmful effects of insecticides on entomopathogenic fungi in integrated pest management programmes (Silva *et al.*, 2013). Hence, the present investigation was carried out to study the interaction of entomopathogenic fungi, *M. anisopliae*, and insecticides under laboratory conditions.

MATERIAL AND METHODS

The investigation of compatibility of *Metarhizium anisopliae* with insecticides was carried out in the Insect pathology laboratory, Department of Agriculture Entomology, Tamil Nadu Agricultural University, Coimbatore, during 2020-2021.

Source of fungal isolate

Metarhizium anisopliae (TNAU-MA-GDU) was found to be highly virulent against Fall armyworm (FAW), *Spodoptera frugiperda* (J. E. Smith) was tested for its compatibility with the insecticides recommended for the management of FAW under laboratory conditions.

Food poison technique

The effect of insecticides on *M. anisopliae* (TNAU-MA-GDU) was assessed by following the poisoned food technique adopted by Neves *et al.*, (2001). The insecticides at required concentrations (0.1X, X, 2X, 5X, and 10X) were added to the sterilized Potato dextrose agar (PDA) medium before solidification and poured into the Petri plates (90 mm diameter) after proper agitation and allowed to solidify. Using a sterilized cork borer, 5mm discs were cut from 7 days old selected EPF and transferred to the center of each plate containing poisoned medium.

*Corresponding author's e-mail: kiruthidg.95@gmail.com

Details of insecticides and their doses used in compatibility studies

Common Name	Trade Name	Dose (X) (ml L ⁻¹ / g L ⁻¹)				
		0.1X	X	2X	5X	10X
Chlorantraniliprole	Coragen 18.5 SC	0.04	0.4	0.8	2.0	4
Flubendiamide	Fame 480 SC	0.05	0.5	1.0	2.5	5
Azadirachtin	Neem 1500	0.5	5.0	10	25	50
Emamectin benzoate	Proclaim 5 SG	0.04	0.4	0.8	2.0	4
Novaluron	Rimon 10 EC	0.15	1.5	3.0	7.5	15
Spinetoram	Delegate 11.70 SC	0.05	0.5	1.0	2.5	5

The untreated check was maintained by placing a fungal disc in a medium without poison for comparison. The plates were incubated at $25 \pm 1^\circ\text{C}$ for 15 days to allow maximum fungal growth. Each treatment consisted of three replications. The diameter of the fungal culture in each plate was measured on the 15th day using a ruler (average of two measurements made at right angles). Fully sporulated cultures were used to determine the spore count using Neubauer haemocytometer. The observations on vegetative growth and sporulation were transformed relative to the control (100) and the product's toxicity (T) value was calculated as per the following formula.

$$T = [20(\text{VG}) + 80(\text{SP})] / 100$$

VG - Vegetative growth; SP - Sporulation

Where, T: 0 to 30 = very toxic; 31 to 45 = toxic; 46 to 60 = moderately toxic; > 60 = compatible.

Effect of insecticides on biomass production of *M. anisopliae* (TNAU-MA-GDU)

For assessment of biomass (dry weight of mycelium), potato dextrose broths (PDB) (media without agar) were prepared. Test concentrations of the insecticides (0.1X, X, 2X, 5X, and 10X) were mixed with 100 ml of sterilized PDB broth separately and a five mm mycelial disc of 10 days old selected fungal culture was inoculated to the broth with help of cork-borer and the flasks were incubated at $25 \pm 1^\circ\text{C}$. The broth culture after 15 days of fungus seeding was filtered through muslin cloth and the mat collected on pre-weighed filter paper was dried at 105°C for 12 h and then reweighed. An untreated check was maintained by placing the fungal disc in PDB broth without poison for comparison. Each treatment was replicated three times. The difference between the final and initial weight was considered as the dry weight of mycelium (Akbar et al., 2012).

Data Analysis

All the experiments were conducted under Completely Randomized Block Design (CRBD). The data obtained in experiments were transformed to square root (X+0.5) transformation and the analysis of variance in different experiments was carried out in AGRES and the means were separated by the least significant difference (LSD) available in the package.

Results and Discussion

Effect of insecticides on vegetative growth and sporulation of *M. anisopliae* (TNAU-MA-GDU)

All the tested insecticides viz., chlorantraniliprole, flubendiamide, azadirachtin, emamectin benzoate, novaluron and spinetoram tested showed significant variation in the vegetative growth and sporulation of *M. anisopliae* (TNAU-MA-GDU) (Table 1 and Figure 1). Based on the product toxicity "T" value, the insecticides were classified into four different categories viz., very toxic, toxic, moderately toxic and compatible. The isolate, *M. anisopliae* (TNAU-MA-GDU) was found to be relatively compatible with all the tested insecticides except Azadirachtin at 0.1 x and x doses based on the T value. The highest vegetative growth (89.0, 88.4, 86.9, 82.5, and 78.8 mm respectively) and spore count (13.3 , 12.8 , 10.4 , 8.7 , and 5.8×10^7 spores mL⁻¹ respectively) of TNAU-MA-GDU isolate was recorded in flubendiamide at 0.1x, x, 2x, 5x and 10x doses followed by spinetoram that recorded the maximum vegetative growth (89.90, 88.1, 86.2, 81.4 and 76.6 mm) and spore count (12.2 , 11.9 , 9.8 , 8.2 and 5.5×10^7 spores mL⁻¹).

All the other insecticides ranged from moderately toxic to toxic based on the "T" value at both twofold higher (2x) and fivefold higher (5x) doses. At a tenfold higher dose (10x), flubendiamide and spinetoram were found to be moderately toxic with a vegetative growth of 48.0 and 46.3 mm and spore count of 5.8 and 5.5×10^7 spores mL⁻¹ whereas, emamectin

benzoate was found to be toxic with vegetative growth and spore count of 32.8 mm and 4.1×10^7 spores mL^{-1} . Azadirachtin was highly toxic to TNAU-MA-GDU isolate at all the test doses with a T value ranging from 43.4 to 4.00.

Effect of insecticides on biomass production of *M. anisopliae* (TNAU-MA-GDU)

The influence of different insecticides on biomass production of *M. anisopliae* isolate (TNAU-MA-GDU) was assessed using PD broth and the results showed that there was a significant difference in biomass production among the insecticides amended broth (Table 2 and Figure 2). The maximum dry weight of the mycelium was recorded in flubendamide treated broth at all the five doses (5.22, 4.97, 4.76, 4.48 and 3.17 g respectively) next to control. This was followed by spinetoram with a biomass production of 5.07, 4.82, 4.51, 4.25, and 3.04 g at a lower dose (0.1x), recommended dose (x), twofold (2x), fivefold (5x), tenfold higher dose (10x), respectively and emamectin benzoate recorded 4.77, 4.48, 4.23, 4.07 and 2.92 g in all the five test doses, respectively. Chlorantraniliprole and novaluron recorded the dry weight ranging from 4.46 to 2.18 and 4.18 to 1.91 g at lower to higher doses. However, azadirachtin recorded the least dry weight (0.10 to 3.34 g) than other insecticides with zero growth at 10x dose.

Current investigations on the interaction effect of *M. anisopliae* (TNAU-MA-GDU) with insecticides indicated that flubendiamide, spinetoram, emamectin benzoate, chlorantraniliprole, and novaluron had synergistic activity at lower and field doses with less effect on vegetative growth, sporulation and biomass production, whereas, azadirachtin had antagonistic effect at all test doses. In contrast, Amutha and Banu, (2012) reported that neem-based formulation, eco neem (Azadirachtin 3%) was harmless to *M. anisopliae*. The contradictions could be due to the quality of neem as influenced by the extraction method and fungus strains used in the study. Similar to present study, the antagonistic effect of Azadirachtin on *M. anisopliae* was also reported by Aguda *et al.* (1986), Hirose *et al.* (2001) and Kumar *et al.* (2008).

Earlier, Akbar *et al.*, (2012) also reported that spinetoram and emamectin benzoate were compatible at tested field doses with minimum inhibition on the growth and sporulation of *M. anisopliae* isolates (M2 and M11). In another study, the insecticides viz., emamectin benzoate, chlorantraniliprole, and novaluron were found to be safer to entomopathogenic fungi with good colony growth and germination percentage (Joshi *et al.*, 2018). Similarly, Parjane *et al.*, (2018) investigated the potential inhibitory effects of 19 pesticides on

the growth of *M. anisopliae* and found that the highest vegetative growth of the fungus was observed in flubendamide, chlorantraniliprole and emamectin benzoate which is in agreement with the present results. Another study made by Tekam *et al.* (2018) also showed that new generation insecticides viz., emamectin benzoate, and flubendiamide were compatible with *M. anisopliae* with least inhibition of 35.24, 37.11, and 39.50 percentage. Similarly, Matcha *et al.* (2021) also found that *M. rileyi* was compatible with chlorantraniliprole, flubendamide, spinetoram, and emamectin benzoate with a spore yield of 2.7, 1.17, 0.9 and 0.26×10^8 spores mL^{-1} which is in line with the present findings




Figure 1. Effect of insecticides on vegetative growth and sporulation of *M. anisopliae* (TNAU-MA-GDU)




Figure 2. Effect of insecticides on biomass production of *M. anisopliae* (TNAU-MA-GDU)

Table 1. Compatibility of *M. anisopliae* (TNAU-MA-GDU) with insecticides

Treatment#	0.1 X			X			2X			5X			10X		
	VG	SP	T Value*	VG	SP	T Value*	VG	SP	T Value*	VG	SP	T Value*	VG	SP	T Value*
Chlorantraniliprole	87.3 (97.0)	10.6 (69.7)	75.2 (D)	85.5 (95.0)	10.1 (66.5)	72.2 (D)	81.5 (90.6)	8.9 (58.6)	64.9 (D)	52.1 (57.9)	4.3 (28.3)	34.2 (B)	45.2 (50.2)	1.1 (7.2)	15.8 (A)
Flubendiamide	89.0 (98.8)	13.3 (87.5)	89.8 (D)	88.4 (98.2)	12.8 (84.2)	87.0 (D)	86.9 (96.6)	10.4 (68.4)	74.1 (D)	82.5 (91.7)	8.7 (57.2)	64.1 (D)	78.8 (87.6)	5.8 (38.2)	48.0 (C)
Azadirachtin	65.2 (72.4)	5.5 (36.2)	43.4 (B)	64.0 (71.1)	5.2 (34.2)	41.6 (B)	59.0 (65.6)	4.8 (31.6)	38.4 (B)	35.2 (39.1)	0.2 (1.3)	8.9 (A)	18.2 (20.2)	0.0 (0.0)	4.0 (A)
Emamectin benzoate	86.0 (95.5)	11.9 (78.3)	81.7 (D)	85.2 (94.7)	11.2 (73.7)	77.9 (D)	83.0 (92.2)	9.1 (59.9)	66.3 (D)	56.9 (63.2)	7.9 (51.9)	54.2 (C)	50.7 (56.3)	4.1 (26.9)	32.8 (B)
Novaluron	86.6 (96.2)	9.5 (62.5)	69.2 (D)	84.0 (93.3)	9.1 (59.9)	66.6 (D)	80.3 (89.2)	7.8 (51.3)	58.9 (C)	49.3 (54.8)	3.9 (25.7)	31.5 (B)	42.1 (46.8)	0.8 (5.3)	13.6 (A)
Spinetoram	89.0 (98.8)	12.2 (80.3)	83.9 (D)	88.1 (97.9)	11.9 (78.3)	82.2 (D)	86.2 (95.8)	9.8 (64.5)	70.7 (D)	81.4 (90.4)	8.2 (53.9)	61.2 (D)	76.6 (86.6)	5.5 (36.1)	46.3 (C)
Untreated control	90.0 (100)	15.2 (100)	-	90.0 (100)	15.2 (100)	-	90.0 (100)	15.2 (100)	-	90.0 (100)	15.2 (100)	-	90.0 (100)	15.2 (100)	-

X- Recommended field dose (Chlorantraniliprole- 0.4 ml L⁻¹; Flubendiamide- 0.5 ml L⁻¹; Azadirachtin- 5 ml L⁻¹; Emamectin benzoate- 0.4 g L⁻¹; Novaluron- 1.5 ml L⁻¹; Spinetoram- 0.5 ml L⁻¹). #Mean of three replications
 VG- Vegetative growth (mm), SP- Sporulation ($\times 10^7$ spores mL⁻¹)
 Values in parentheses are values relative to control (100 %)
 *Classification based on formula T= (20VG+80SP)/100; A- Very Toxic, B- Toxic, C- Moderately toxic, D- Compatible

Table 2. Effect of insecticides on biomass production of *M. anisopliae* (TNAU- MA-GDU)

Treatment#	Biomass* (g)				
	0.1 X	X	2X	5X	10X
Chlorantraniliprole	4.46 (2.11) ^d	4.15 (2.04) ^d	3.99 (2.00) ^e	3.27 (1.81) ^e	2.18 (1.48) ^d
Flubendiamide	5.22 (2.28) ^{ab}	4.97 (2.23) ^b	4.76 (2.18) ^b	4.48 (2.12) ^b	3.17 (1.78) ^b
Azadirachtin	3.34 (1.83) ^f	3.10 (1.76) ^f	2.92 (1.71) ^g	0.10 (0.32) ^g	0.00 (0.71s) ^f
Emamectin benzoate	4.77 (2.18) ^c	4.48 (2.12) ^c	4.23 (2.06) ^d	4.07 (2.02) ^d	2.92 (1.71) ^c
Novaluron	4.18 (2.04) ^e	3.94 (1.98) ^e	3.44 (1.85) ^f	2.98 (1.73) ^f	1.91 (1.38) ^e
Spinetoram	5.07 (2.25) ^b	4.82 (2.20) ^b	4.51 (2.12) ^c	4.25 (2.06) ^c	3.04 (1.74) ^{bc}
Untreated control	5.49 (2.34) ^a	5.49 (2.34) ^a	5.49 (2.34) ^a	5.49 (2.34) ^a	5.49 (2.34) ^a
SEd	0.0278	0.0185	0.0150	0.0188	0.0185
CD (0.05)	0.0596	0.0397	0.0321	0.0403	0.0397

X- Recommended field dose (Chlorantraniliprole- 0.4 ml L⁻¹; Flubendiamide- 0.5 ml L⁻¹; Azadirachtin- 5 ml L⁻¹; Emamectin benzoate- 0.4 g L⁻¹; Novaluron- 1.5 ml L⁻¹; Spinetoram- 0.5 ml L⁻¹).
 *Figures in the parentheses are square root ($\sqrt{x+0.5}$) transformed values.
 In a column, means followed by a common letter are not significantly different at the 5% level by LSD.
 #Mean of three replications



CONCLUSION

The isolate, *M. anisopliae* (TNAU-MA-GDU) was compatible with flubendiamide, spinetoram, emamecton benzoate, chlorantraniliprole, and novaluron at field doses and hence can be combined for the management of Fall armyworm under field conditions. Green labeled insecticides, azadirachtin was toxic at all test doses and inhibited the growth and sporulation of *M. anisopliae* (TNAU-MA-GDU). Hence, it demonstrated that insecticides except azadirachtin had a synergistic effect on *M. anisopliae* at field doses under laboratory conditions. It needs to be further validated through field experiments.

Funding

Not applicable

Acknowledgment

The authors gratefully acknowledge the facilities provided under the GoTN – FAW project for carrying out the above research.

Ethics statement

Not applicable

Originality and plagiarism

The authors assure that the research work submitted here is original and not subjected to any plagiarized content.

Consent for publication

All the authors agreed to publish the content.

Competing interests

There was no conflict of interest in the publication of this content

Data availability

Not applicable

Author contributions

Idea conceptualization- SJ, Guidance- SJ, NS, MM, US, DU, Experiments- GK, Writing original draft- GK, Writing- reviewing & editing- SJ.

REFERENCES

Akbar, S., Freed, S., Hameed, A., Gul, H. T., Akmal, M., Malik, M. N., Naeem, M. and M. B. Khan. 2012. Compatibility of *Metarhizium anisopliae* with different insecticides and fungicides. *Afr. J. Microbiol. Res.*, **6(17)**: 3956-3962.

Aguda, R. M., Rombach, M. C., and B. M. Shepard, 1986. Effect of neem oil on germination and sporulation of the entomogenous fungus *Metarhizium anisopliae* (Philippines). *International Rice Research Newsletter.*, **1(4)**: 34-35.

Asi, M. R., Bashir, M. H., Afzal, M., Ashfaq, M., and S. T. Sahi. 2010. Compatibility of entomopathogenic fungi, *Metarhizium anisopliae* and *Paecilomyces fumosoroseus* with selective insecticides. *Pak. J. Bot.*, **42(6)**: 4207-4214.

Hirose, E., Neves, P. M., Zequi, J. A., Martins, L. H., Peralta, C. H., and A. Moino Jr. 2001. Effect of biofertilizers and neem oil on the entomopathogenic fungi *Beauveria bassiana* (Bals.) Vuill. and *Metarhizium anisopliae* (Metsch.) Sorok. *Braz. Arch. Biol. Technol.*, **44**: 419-423.

Joshi, M., Gaur, N., and R. Pandey. 2018. Compatibility of entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* with selective pesticides. *J. Entomol. Zool. Stud.*, **6(4)**: 867-872.

Kumar, A. M., Reddy, K. N., and R. Sreevathsa. 2008. Influence of pesticides, plant oils and antagonistic entomopathogenic fungus, *Metarhizium anisopliae* (Metsch.) Sorokin. *Pest Technol.*, **2(1)**: 28-31.

Matcha, N., Duraimurugan, P., and A. K. Bhowmick. 2021. Effect of insecticides and fungicides on growth and sporulation of *Metarhizium rileyi* (Farlow) Samson. *J. Pharm. Innov.*, **10(7)**: 1444-1447.

Neves, P. M., Hirose, E., Tchujo, P. T., and J. R. Moino. 2001. Compatibility of entomopathogenic fungi with neonicotinoid insecticides. *Neotrop. Entomol.*, **30**: 263-268.

Oliveira, C. N. D., Neves, P. M., and L. S. Kawazoe. 2003. Compatibility between the entomopathogenic fungus *Beauveria bassiana* and insecticides used in coffee plantations. *Sci. Agric.*, **60**: 663-667.

Parjane, N. V., Kabre, G. B., Mahale, A. S., Shejale, B. T., and S. A. Nirgude. 2020. Compatibility of pesticides with *Metarhizium anisopliae*. *J. Entomol. Zool. Stud.*, **8(5)**: 633-636.

Purwar, J. P., and G. C. Sachan. 2006. Synergistic effect of entomogenous fungi on some insecticides against Bihar hairy caterpillar *Spilarctia obliqua* (Lepidoptera: Arctiidae). *Microbiol. Res.*, **161(1)**: 38-42.

Sain, S. K., Monga, D., Kumar, R., Nagrale, D. T., Hiremani, N. S., and S. Kranthi. 2019. Compatibility of entomopathogenic fungi with insecticides and their efficacy for IPM of *Bemisia tabaci* in cotton. *J. Pestic. Sci.*, **44(2)**: 97-105.

Schumacher, V., and H. M. Poehling. 2012. In vitro effect of pesticides on the germination, vegetative growth, and conidial production of two strains of *Metarhizium anisopliae*. *Fungal Biol.*, **116(1)**: 121-132.

Silva, R. A. D., Quintela, E. D., Mascarin, G. M., Barrigossi, J. A. F., and L. M. Lião. 2013. Compatibility of conventional agrochemicals used in rice crops with the entomopathogenic fungus *Metarhizium anisopliae*. *Sci. Agric.*, **70**: 152-160.

Tekam, K. D., Kelwatkar, N. M., and S. B. Das. 2018. Compatibility of *Metarhizium anisopliae* with new generation insecticide in vitro condition. *J. Entomol. Zool. Stud.*, **6(6)**: 887-890.